



WELCOME

A guide to **getting started** with **Evrysdi**



What is Evrysdi?

Evrysdi is a prescription medicine used to treat spinal muscular atrophy (SMA) in children and adults.

Please see additional Important Safety Information and full Prescribing Information.



HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Welcome
to Evrysdi!

This guide is
here to help you get
started with treatment.

Find answers to your
questions and helpful
resources by clicking
the buttons to the left.



Important Safety Information

Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant, as Evrysdi may harm your unborn baby. Ask your healthcare provider for advice before taking this medicine

Please see additional Important Safety Information and full Prescribing Information.

 **Evrysdi**[®]
risdiplam 60 mg
powder for oral solution

HOME

**WHERE
TO START**

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Where to start: your **local contact**

The PAL (Partnership and Access Liaison) team are local points of contact at Genentech to answer your questions and offer support and guidance.

A PAL can:

- ✓ **Provide** in-person or virtual support
- ✓ **Answer** questions about Evrysdi
- ✓ **Explain** insurance coverage and potential financial support options
- ✓ **Refer** you to helpful resources

Important Safety Information (continued)

Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:

- are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi

Please see additional Important Safety Information and full Prescribing Information.

Don't hesitate
to reach out!

Get connected at
Evrysdi.com/PAL
(833) 387-9734
(Monday-Friday,
9 AM to 8 PM ET)

Evrysdi[®]
risdiplam 60 mg
powder for oral solution

Where to start: a few **things to note**

HOME

**WHERE
TO START**

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information



- ✓ Evrysdi is delivered directly to your door
- ✓ A specialty pharmacy* will call you to schedule a time for your delivery
- ✓ Each delivery includes all the medicine and supplies you'll need for ~1 month†
- ✓ You will need to store your Evrysdi in the refrigerator in the original amber bottle to protect it from light
- ✓ Each shipment includes up to 3 bottles of liquid medicine

*Specialty pharmacies are not part of Genentech and maintain independence in their operations and in their role as a healthcare provider.

†The exact amount of Evrysdi you receive is based on your dose and how much your insurance plan allows to be delivered at one time.

Important Safety Information (continued)

Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:

- are a woman who can become pregnant:
 - **Pregnancy Registry.** Talk to your healthcare provider right away if you become pregnant while taking Evrysdi. Ask about registering with the Evrysdi Pregnancy Registry, which was created to collect information about your health and your baby's health. Your healthcare provider can enroll you in this registry by calling 1-833-760-1098 or visiting www.evrysdipregnancyregistry.com


Please see additional Important Safety Information and full Prescribing Information.



Where to start: a few **things to note**



- ✓ Plan to take Evrysdi by mouth or feeding tube once a day after a meal*
- ✓ It can be helpful to set reminders on your phone or calendar



- ✓ Your healthcare provider will determine your dose based on your age and weight



- ✓ Financial support may be available to you
- ✓ You can find more information [here](#)

These are just a few elements to keep in mind.
For detailed **Instructions for Use**, click [here](#).

*Evrysdi must be taken within 5 minutes of when it is drawn up into the syringe.

Important Safety Information (continued)

Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:

- are an adult male. Evrysdi may affect a man's ability to have children (fertility). Ask a healthcare provider for advice before taking this medicine

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

**WHERE
TO START**

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information



Where to start: when you have **questions**

HOME

**WHERE
TO START**

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

GO TO...

✓ **Your healthcare provider**

✓ **The specialty pharmacy**
(the company that will call you to schedule your delivery)

✓ **MySMA Support* or a PAL**
(Partnership and Access Liaison)
(833) 387-9734, Monday-Friday
9 AM to 8 PM ET

FOR QUESTIONS ABOUT...

What to expect when starting treatment, your dose, potential side effects, your results on Evrysdi

Your delivery of Evrysdi

Potential financial support options, storing or taking your medicine, planning for travel, helpful resources — you can also ask your healthcare provider or specialty pharmacy for help with these elements

*Enrollment in MySMA Support through the Evrysdi Start Form is mandatory to receive assistance through the program. Participation in MySMA Support is not necessary to receive treatment with Evrysdi.

Important Safety Information (continued)

Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:

- are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby

Please see additional Important Safety Information and full Prescribing Information.



HOME

WHERE
TO START

**STORING
EVRYSDI**

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Storing Evrysdi: your **delivery**

Evrysdi is delivered directly to your door by a specialty pharmacy.* Evrysdi is not available at retail pharmacies

- You will receive a call from the specialty pharmacy to welcome you and schedule a delivery time that works best for you; your preferred delivery time will be confirmed each month
- Each delivery includes an ~1-month supply[†] of medication and materials

*Specialty pharmacies are not part of Genentech and maintain independence in their operations and in their role as a healthcare provider.

[†]The exact amount of Evrysdi you receive is based on your dose and how much your insurance plan allows to be delivered at one time.

If you have **questions about your delivery**,
call your specialty pharmacy.

Important Safety Information (continued)

Tell your healthcare provider about all the medicines you take.

Please see additional Important Safety Information and full Prescribing Information.



Storing Evrydsi: your **delivery**

HOME

WHERE
TO START

**STORING
EVRYSDI**


TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Each package includes:



Your shipment will also include full [Instructions for Use](#) and [Patient Information](#).

The specialty pharmacy will select the appropriate oral syringe (1 mL, 6 mL, or 12 mL) based on the dose prescribed by your doctor and will include 1 or 2 in your shipment.

Important Safety Information (continued)

You should receive Evrydsi from the pharmacy as a liquid. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



Storing Evrydsi: at **home** or when **traveling**

HOME

WHERE
TO START

**STORING
EVRYSDI**

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Storing your medicine

- Always store Evrydsi upright in the refrigerator (36°F to 46°F) in the original amber bottle to protect it from light. Do not freeze Evrydsi or put the medicine in a different container
- If refrigeration is not available, Evrydsi can be kept at room temperature up to 104°F for a combined total of 5 days
- It can be removed from, and returned to, a refrigerator
- The total combined time out of refrigeration should not exceed 5 days
- Any portion of Evrydsi that has been kept above 104°F should be discarded
- Always store your medication in a safe place out of the reach of children
- Discard any unused medication after 64 days. Be sure to check the “Discard After” date written on the bottle label, and carefully read the **Instructions for Use** that comes with your Evrydsi
- To receive a travel cooler to help store your Evrydsi, please contact a PAL at **(833) 387-9734**, Monday-Friday, 9 AM to 8 PM ET

Important Safety Information (continued)

Avoid getting Evrydsi on your skin or in your eyes. If Evrydsi gets on your skin, wash the area with soap and water. If Evrydsi gets in your eyes, rinse your eyes with water.

Please see additional Important Safety Information and full Prescribing Information.



HOME

WHERE
TO START

STORING
EVRYSDI

**TAKING
EVRYSDI**

SUPPORT

FAQs

Important
Safety
Information

Taking Evrysdi: your **dose**

Your healthcare provider determined your dose based on your age and weight. Always take your dose exactly as prescribed.

Be sure to tell your healthcare provider if you or the person in your care is taking:

- Any other medications, vitamins, or dietary supplements, including any stool softeners
- Additional fiber that may impact bowel movements, as these may need to be adjusted prior to starting Evrysdi

If you have **questions about your dose**, speak with your healthcare provider.

Important Safety Information (continued)

The most common side effects of Evrysdi include:

- For later-onset SMA: fever, diarrhea, rash

Please see additional Important Safety Information and full Prescribing Information.



Taking Evrysdi: **preparing** your medicine

HOME

WHERE
TO START

STORING
EVRYSDI

**TAKING
EVRYSDI**

SUPPORT

FAQs

Important
Safety
Information

Before preparing your first dose



Please read the full [Patient Information](#) and [Instructions for Use](#) that come with your prescription and discuss with your healthcare provider before taking the first dose.

Important Safety Information (continued)


The most common side effects of Evrysdi include:

- For infantile-onset SMA: fever; diarrhea; rash; runny nose, sneezing, and sore throat (upper respiratory infection); lung infection (lower respiratory infection); constipation; vomiting; cough

These are not all of the possible side effects of Evrysdi. For more information on the risk and benefits profile of Evrysdi, ask your healthcare provider or pharmacist.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).

To start, take a clean oral syringe out of the box



Your package comes with 1 or 2 reusable oral syringe(s) for taking or giving Evrysdi. **Always take or give the dose exactly as prescribed by your healthcare provider, and administer with the reusable syringes that come with your prescription (they help protect Evrysdi from light).**



Taking Evrydsi: **preparing** your medicine

HOME

WHERE
TO START

STORING
EVRYSDI


**TAKING
EVRYSDI**

SUPPORT


FAQs

Important
Safety
Information

Next, fill the oral syringe with medication



Take off the child-resistant cap by pushing it down and twisting to the left. Do not throw away the cap.



Push the plunger of the syringe all the way down to remove any air inside.



Place the bottle upright on a flat surface, such as a table or counter. Insert the syringe tip into the bottle adapter.

**If you
spill
Evrydsi**

Dry the area with a dry paper towel and then clean with water. Throw away the paper towel in the trash and wash your hands well with soap and water. You do not need to wear gloves.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



Taking Evrydsi: **preparing** your medicine

HOME

WHERE
TO START

STORING
EVRYSDI


**TAKING
EVRYSDI**

SUPPORT


FAQs

Important
Safety
Information


Fill the oral syringe with medication (continued)



Carefully turn the bottle upside down, with the syringe tip firmly inserted into the bottle adapter.



Slowly pull back on the plunger to withdraw the prescribed dose. The top of the black stopper should line up with the mL marking that corresponds to the prescribed dose volume.



After the correct dose is withdrawn, **continue to hold the plunger in place to keep it from moving**. Leave the syringe in the bottle adapter and turn the bottle upright again. Return the bottle to the flat surface and gently remove the syringe by pulling straight up while holding the plunger in place.

**If you
spill
Evrydsi**

Dry the area with a dry paper towel and then clean with water. Throw away the paper towel in the trash and wash your hands well with soap and water. You do not need to wear gloves.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).

 **Evrydsi**[®]
risdiplam 60 mg
powder for oral solution

Taking Evrydsi: **preparing** your medicine

HOME

WHERE
TO START

STORING
EVRYSDI


**TAKING
EVRYSDI**

SUPPORT


FAQs

Important
Safety
Information

Fill the oral syringe with medication (continued)



Hold the syringe with the tip pointing up. Check the syringe to ensure there aren't any air bubbles in the medication. If you see bubbles (especially larger ones), you may not have the correct amount of medication. **If bubbles occur, put the syringe tip firmly into the bottle adapter and use the plunger to put the medication back in the bottle, then repeat steps 4-7.**



Return the cap to the bottle. Turn the cap to the right and make sure it's on tight before returning the bottle to the refrigerator.

**Give or take
Evrydsi right
away after it's
drawn into the
syringe**

If it is not taken within **5 minutes**, do not take that dose, empty the syringe by throwing away the unused medication in the household trash (but do not throw away the syringe), and prepare a new dose.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).


 **Evrydsi**[®]
risdiplam 60 mg
powder for oral solution

Taking Evrysdi: by **mouth**

You or the person in your care should sit upright while taking Evrysdi.



Place the tip of the oral syringe inside the mouth, touching the inside of either cheek. **To avoid choking, do not put the syringe toward the back of the mouth.** Slowly push the plunger all the way down until all of the medication is emptied from the syringe. **Pushing the plunger too quickly may cause choking.**



Check the syringe to make sure there is no Evrysdi left inside.



You or the person in your care should drink some water (about 1 tablespoon) after swallowing the medication to ensure all the medication is swallowed.

When to take Evrysdi

You or the person in your care should take Evrysdi at the same time every day.
Evrysdi should be taken after a meal. It should not be mixed directly into foods or other liquids.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE TO START

STORING EVRYSDI

TAKING EVRYSDI

SUPPORT

FAQs

Important Safety Information

Taking Evrysdi: by **feeding tube**

HOME

WHERE
TO START


STORING
EVRYSDI

**TAKING
EVRYSDI**


SUPPORT

FAQs


Important
Safety
Information



Place the tip of the syringe into the G- or NG-tube.
Slowly push the plunger all the way down until the syringe is empty.



Be sure to check the syringe to ensure all Evrysdi has been given.



Flush the feeding tube with water (about 2 to 4 teaspoons) right after giving the prescribed dose.

You can
find a video
demonstrating
how to take
Evrysdi [here](#)

G-tube stands for gastrostomy tube; NG-tube stands for nasogastric tube.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



Taking Evrysdi: **what to do** if...

IMPORTANT REMINDERS

If the medication is spit up or not fully swallowed

- Do not give any more medication that day
- Wait until the next day to give the next dose at the usual dosing time

If a dose is missed

- The missed dose should be taken as soon as possible, but no more than 6 hours after the usual dosing time
- If more than 6 hours have passed, skip the missed dose altogether and take or give the next dose the following day



HOME

WHERE
TO START

STORING
EVRYSDI

**TAKING
EVRYSDI**

SUPPORT

FAQs


Important
Safety
Information

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).

 **Evrysdi**[®]
risdiplam 60 mg
powder for oral solution

Taking Evrysdi: **cleaning** up


Clean the oral syringe for reuse



1 Separate the plunger from the syringe by pulling the plunger away from the syringe tip until it comes out of the syringe completely.



2 Rinse both pieces thoroughly with water.



3 Once clean, let them dry on a clean surface.



4 Be sure to wash your hands with soap and water after cleaning the syringe.

After the oral syringe barrel and plunger are dry, put the plunger back into the oral syringe barrel, and store the syringe with your medicine.

Always clean the syringe and plunger by hand, and never in the dishwasher

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE TO START

STORING EVRYS DI

TAKING EVRYS DI

SUPPORT

FAQs

Important Safety Information

Taking Evrysdi: **cleaning** up

How to dispose of Evrysdi

- Follow your local guidelines for proper disposal of medication. Visit www.fda.gov/drugs/ensuring-safe-use-medicine/safe-disposal-medicines or contact your healthcare provider for more information

If there is something wrong with your delivery

- If you receive an incomplete order, a broken bottle, or medication in powder form, do not take the medication you received, and contact your specialty pharmacy immediately
- Contact your specialty pharmacy with questions about shipping and packaging

If you notice leakage on any bottle, dry off the bottle with a dry paper towel, then rinse the closed bottle with water. Store the bottle out of sight and out of reach of children. Call Genentech Product Complaints at **1-800-334-0290** to coordinate your free replacement and receive further instructions.

If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE
TO START

STORING
EVRYSDI

**TAKING
EVRYSDI**

SUPPORT

FAQs

Important
Safety
Information

Taking Evrysdi: important **reminders**

DO

- ✓ Do ask your healthcare provider how to use the reusable oral syringe and how to measure your prescribed daily dose
- ✓ Do always use the oral syringes provided and contact your specialty pharmacy immediately if you receive oral syringes that are not identical in size
- ✓ Do take the dose of Evrysdi that your healthcare provider has prescribed to you or the person in your care
- ✓ Do take the dose at the same time every day after a meal
- ✓ Do store medication upright in the original amber bottle and close the bottle cap completely after using
- ✓ Do keep Evrysdi cool (36°F to 46°F). Evrysdi should be stored in the refrigerator. Always store medication in a safe place out of the reach of children
- ✓ Do check the expiration date so you don't use expired medication (Evrysdi expires 64 days after being prepared by a pharmacist)
- ✓ Do ask your specialty pharmacy for the **Discard After** date if it is not written on the bottle label

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE
TO START

STORING
EVRYSDI

**TAKING
EVRYSDI**

SUPPORT

FAQs

Important
Safety
Information

Taking Evrysdi: important **reminders**

DO NOT

- ✗ Do not use Evrysdi if the bottle adapter is not in the bottle. Contact your specialty pharmacy immediately if you receive a bottle without an adapter
- ✗ Do not use Evrysdi if it is in powder form, or if the bottle is damaged. Contact your specialty pharmacy immediately if you receive a bottle containing Evrysdi in powder form
- ✗ Do not get Evrysdi on your skin or in your eyes. If it gets on the skin, wash with soap and water. If it gets in your eyes, flush with water
- ✗ Do not mix Evrysdi into foods, liquids, formula, or milk
- ✗ Do not use Evrysdi if the bottle or oral syringes are damaged
- ✗ Do not freeze Evrysdi
- ✗ Do not put the medication in a different container from the one it comes in
- ✗ Do not take a partial dose, even if that is all you have left in a bottle
- ✗ Do not store Evrysdi doses in the oral syringe — discard any doses that have been in the syringe for more than 5 minutes and prepare a new dose

If you have questions about how to take Evrysdi, contact your healthcare provider. A PAL may also be able to answer questions about Evrysdi; to connect with a PAL, call **(833) 387-9734**, Monday-Friday, 9 AM to 8 PM ET.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE
TO START

STORING
EVRYSDI

**TAKING
EVRYSDI**

SUPPORT

FAQs

Important
Safety
Information

Support: we're **here for you**

MySMA Support* is a service from Genentech that can help provide information about Evrysdi.

The team is available to:



answer your questions



provide information about Evrysdi, including how to take it



help you understand your insurance coverage



refer you to appropriate Evrysdi financial assistance options



coordinate the preparation and delivery of your medicine with the specialty pharmacy

To contact the MySMA Support team:

Visit [Evrydsi.com/PAL](https://www.evrydsi.com/PAL)

Call **(833) 387-9734** (Monday-Friday, 9 AM to 8 PM ET)

MySMA Support, including the PAL, does not provide medical advice and is not a substitute for your medical team. Your healthcare provider should always be your main resource for any questions about your health and medical care.

*Enrollment in MySMA Support through the Evrysdi Start Form is mandatory to receive assistance through the program. Participation in MySMA Support is not necessary to receive treatment with Evrysdi.

Please see additional Important Safety Information and full Prescribing Information.



HOME

WHERE TO START

STORING EVRYSDI

TAKING EVRYSDI

SUPPORT

FAQs

Important Safety Information

Support: we're **here for you**

HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

MEET THE TEAM

Your primary contact: **PALs**

Partnership and Access Liaisons (PALs) are the local, main points of contact from Genentech here to support you. Contact a PAL for answers to questions about Evrysdi (including how it works, how to take it, what results were seen in clinical trials, and Important Safety Information), for help understanding your insurance coverage and potential financial support options, and to be connected to helpful resources.

Additional support: **Case Managers**

While a PAL is your main point of contact, Case Managers partner closely with PALs and healthcare providers to help you understand the health insurance process and identify potential financial support options for Evrysdi.

MySMA Support, including the PAL, does not provide medical advice and is not a substitute for your medical team. Your healthcare provider should always be your main resource for any questions about your health and medical care.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



Support: **financial** options

Your insurance coverage affects the type of financial support you may be eligible for. Depending on your insurance, support options may include one of the following:

Evrysdi Co-pay Program*

You pay as little as \$0 per prescription for Evrysdi if you qualify for the Evrysdi Co-pay program. The program covers the rest, up to \$25,000 per calendar year.

*This Evrysdi Co-pay Program is valid ONLY for patients with commercial insurance who have a valid prescription for a Food and Drug Administration (FDA)-approved indication of a Genentech medication. Patients using Medicare, Medicaid or any other federal or state government program to pay for their medicines are not eligible. Full program terms and conditions can be found [here](#).

Independent co-pay foundations†

If you need help with the co-pay for your Evrysdi, we can refer you to an independent co-pay assistance foundation.

†Independent co-pay assistance foundations have their own rules for eligibility. Genentech has no involvement or influence in independent foundation decision-making or eligibility criteria and does not know if a foundation will be able to help you. We can only refer you to a foundation that supports your disease state. Genentech does not endorse or show preference for any particular foundation. The foundations we refer you to may not be the only ones that might be able to help you.

The Genentech Patient Foundation‡

The Genentech Patient Foundation gives free Evrysdi to people who don't have insurance coverage or who have financial concerns and meet eligibility criteria.

‡If you have health insurance, you should try to get other types of financial assistance, if available. You also need to meet income requirements. If you do not have insurance, or if your insurance does not cover your Genentech medicine, you must meet a different set of income requirements. Genentech reserves the right to modify or discontinue the program at any time and to verify the accuracy of information submitted.

For more information

A detailed guide to Insurance and Financial Assistance Options can be found on our [website](#).

If you have questions about possible financial support options, contact MySMA Support at **(833) 387-9734** (Monday-Friday, 9 AM to 8 PM ET)

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE TO START

STORING EVRYSDI

TAKING EVRYSDI

SUPPORT

FAQs

Important Safety Information

Support: helpful **reminders**

HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

✓ Remembering to take your daily dose

It can be helpful to set daily reminders on your phone or calendar, and to take Evrysdi at the same time each day, so it can become part of your routine.

✓ Support for insurance questions

The MySMA Support team can help navigate insurance and financial assistance questions and coordinate with your healthcare provider and insurance company.* You can reach them Monday-Friday, 9 AM to 8 PM ET at (833) 387-9734. You can also find detailed information on our [website](#).

✓ Planning ahead for renewals

If you and your doctor decide to continue with your Evrysdi prescription, your insurance company may require specific tests every 6 to 12 months to approve the renewal of your Evrysdi prescription. It's a good idea to plan ahead for any appointments you may need in advance. The MySMA Support team can help you determine what may be required.

*You and your doctor are responsible for completing and submitting all required paperwork to your health insurance plan. Genentech cannot guarantee your plan will cover any treatments.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



Support: tracking **your results**

Since everyone's experience with a new treatment can be different, it can be helpful to keep track of your goals and results over time, and to discuss both with your healthcare provider.

Print these pages when starting treatment for a simple way to keep your notes organized.

Why I chose Evrysdi: _____

The skills I have now that are important for me to maintain: _____

My expectations for the first 6 months on Evrysdi: _____

My expectations over time (years): _____

How it's going (how I feel, changes I've noticed):

After 1 month _____

After 4 months _____

After 8 months _____

After 1 year _____

My scores (on tests my doctor may use to measure my progress):

Before starting Evrysdi _____

After 6 months _____

After 1 year _____

After 1.5 years _____

After 2 years _____

**Just
a note!**

The welcome kit you receive in the mail also includes a blank journal you can use to track your results and make notes.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Support: **questions** for your **healthcare provider**

Questions to discuss with my doctor:

- Should I adjust any of the medications I am taking (including vitamins, fiber supplements, laxatives, and prescription medicines)?
- Are there any aspects of my nutrition that we may need to adjust, including my diet or typical meal times? Would it be helpful to talk to a nutritionist?
- What side effects could I experience when starting treatment or as I continue on treatment?
- If I do experience side effects, what should my plan be to address them?
- How should I expect to feel on treatment?
- What results should I expect from treatment?
- How will we measure my progress?
- Are there any tests I may need to plan for to renew my prescription each year?

Additional questions:

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Support: **things you can do** about **common side effects**

Every time you start a new medication, it's important to know about the potential side effects, what to look out for, and steps to take if they happen.

As always, your doctor is the best resource for any questions. It's important to talk to your doctor if you notice any changes, begin any new medication, or make changes in your overall care. Below are some tips that may help if you experience these changes.

If you experience diarrhea:

- It means you have loose, watery, and more-frequent bowel movements
- Some dietary tips you can consider include:
 - Drinking more fluids, such as water, broths, and juices
 - Avoiding juices or foods that have a laxative effect, such as prunes or prune juice
 - Eating semisolid and low-fiber food gradually, such as soda crackers, toast, eggs, rice, or chicken
 - Avoiding dairy products, fatty foods, high-fiber foods, or highly seasoned foods for a few days
- In Evrysdi clinical studies, 17% of adults and children taking Evrysdi experienced diarrhea, with a similar frequency in infants. The majority of cases* (8 out of 10) lasted less than 1 week; 12% lasted longer than 2 weeks
- Talk to your healthcare provider if you are taking other medications, vitamins, or dietary supplements, including any stool softeners or additional fiber that may impact bowel movements

If you experience rash:

- It means you have an area of irritated or swollen skin; it may be itchy and red
- Some tips you can consider include:
 - Using moisturizer or lotions
 - Taking a cool bath
 - Avoiding harsh soaps
 - Using cortisone cream to relieve swelling (if not infected)
 - Avoiding wearing tight clothing

*Cases with a known duration.

Please see additional Important Safety Information and full Prescribing Information.



HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Support: important **phone numbers**

For an easy reference, enter your key contacts in the fields below, then click save.

My PAL:

Specialty Pharmacy:

MySMA Support:

(833) 387-9734

Notes:

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Frequently asked **questions**



Can I get Evrysdi from a retail pharmacy?

Evrysdi is not available at a retail pharmacy, as it requires specific mixing and handling only available at a specialty pharmacy.



If my delivery arrives later than expected, how do I know it's safe for use?

Contact your specialty pharmacy representative right away to confirm your product is safe to take.



What should I do if there is an issue with my order, if any of the contents of my shipment (bottles, oral syringes, etc.) are damaged, or if the seal on my medication is broken?

Contact your specialty pharmacy representative right away.



What should I do if my bottle of Evrysdi is leaking?

If you notice leakage on any bottle, dry off the bottle with a dry paper towel, then rinse the closed bottle with water. Store the bottle out of sight and out of reach of children. Call Genentech Product Complaints at **1-800-334-0290** to coordinate your free replacement and receive further instructions. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water.



What if I have trouble with the packaging?

If you have trouble with the packaging, contact your specialty pharmacy.



Should my product have a temperature tag?

Evrysdi does not require a temperature tag. You should keep Evrysdi refrigerated at 36°F to 46°F at all times. Contact your specialty pharmacy if you have any questions about the correct temperature for your medicine.



Can Evrysdi be kept at room temperature?

If refrigeration is not available, Evrysdi can be kept at room temperature up to 104°F for a combined total of 5 days. It can be removed from, and returned to, a refrigerator. The total combined time out of refrigeration should not exceed 5 days.

Any portion of Evrysdi that has been kept above 104°F should be discarded.

Contact your specialty pharmacy if you have any questions.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Frequently asked **questions**



How long can Evrysdi be stored in the refrigerator?

The medication lasts up to 64 days after being prepared by a pharmacist. An expiration date will be indicated on the bottle of Evrysdi. It may be helpful to mark that date on your calendar. If you receive a bottle without the Discard After date written on the bottle label, contact your specialty pharmacy.



How should I store Evrysdi while traveling?

If you or the person in your care will be away from home, store Evrysdi in an insulated cooler (36°F to 46°F with ice packs), but do not freeze it. Store Evrysdi upright in the original amber bottle to protect it from light. We can send you an Evrysdi travel cooler if you need one. If traveling by plane, your healthcare provider may give you a letter authorizing you to fly with liquid medication. Talk to your healthcare provider about the best way to travel with Evrysdi.



Can I take Evrysdi with food or mix it with liquid outside of the oral syringe?

Evrysdi should be taken after a meal. It should not be mixed directly with any food or liquid (including formula or breast milk) and should only be taken with the approved oral syringe, which has been designed to protect the solution from light.



What do I need to know about nutrition when starting Evrysdi?

Before starting treatment, tell your healthcare provider what foods you or the person in your care generally eats, as well as your typical meal times. Discuss if any part of your routine may need to be adjusted, such as dairy products, fatty foods, high-fiber foods, highly seasoned foods, or foods that have a laxative effect.



What should I do if my child spits up, vomits, or doesn't swallow the whole dose after I give the medication?

Do not give any more medication that day. Wait until the next day to give the next dose at the usual dosing time. If your child continues to spit up or vomit while taking Evrysdi, talk to your healthcare provider.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

Important
Safety
Information

Frequently asked **questions**

✓ **What side effects should I expect when starting Evrysdi?**

Everyone reacts differently when starting a new treatment. Side effects can be impacted by a number of elements, including other medications you may be taking.

Before starting treatment, tell your healthcare provider if you or the person in your care is taking any other medications, vitamins, or dietary supplements, including any stool softeners or additional fiber that may impact bowel movements. Your healthcare provider may wish to adjust these medications prior to starting Evrysdi.

The most common side effects of Evrysdi for later-onset SMA and infantile-onset SMA include fever, diarrhea, and rash. Infantile-onset SMA also included upper respiratory infection, lower respiratory infection, constipation, vomiting, and cough.

In clinical trials, 17% of adults and children and 15% of infants taking Evrysdi experienced diarrhea. The majority (8 out of 10) of cases* of diarrhea lasted less than 1 week and 12% of cases lasted longer than 2 weeks. **See page 28 for some helpful tips if you experience diarrhea, and be sure to speak with your healthcare provider.**

✓ **What if a dose is missed?**

The missed dose should be taken as soon as possible, but no more than 6 hours after the usual dosing time. If more than 6 hours have passed, skip the missed dose altogether and take or give the next dose the following day at your regularly scheduled time.

✓ **Who should I call if I have other questions about Evrysdi?**

Your healthcare provider should always be your main resource for any questions about your health and medical care. You can also contact MySMA Support to connect with a PAL, who can answer questions about Evrysdi, and refer you to helpful resources. PALs can also help you understand your insurance and financial support options. Contact MySMA Support at **(833) 387-9734** (Monday-Friday, 9 AM to 8 PM ET).

*Cases with a known duration.

Please see additional Important Safety Information and full Prescribing Information.



HOME

WHERE
TO START

STORING
EVRYSDI

TAKING
EVRYSDI

SUPPORT

FAQs

**Important
Safety
Information**

Important Safety Information

What is Evrysdi?

Evrysdi is a prescription medicine used to treat spinal muscular atrophy (SMA) in children and adults

Important Safety Information

Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:



- are pregnant or plan to become pregnant, as Evrysdi may harm your unborn baby. Ask your healthcare provider for advice before taking this medicine
- are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi
 - **Pregnancy Registry.** There is a pregnancy registry for women who take EVRYSDI during pregnancy. The purpose of this registry is to collect information about the health of the pregnant woman and her baby. If you are pregnant or become pregnant while receiving EVRYSDI, tell your healthcare provider right away. Talk to your healthcare provider about registering with the EVRYSDI pregnancy Registry. Your healthcare provider can enroll you in this registry or you can enroll by calling 1-833-760-1098 or visiting <https://www.evrysdipregnancyregistry.com>.



- are an adult male. Evrysdi may affect a man's ability to have children (fertility). Ask a healthcare provider for advice before taking this medicine



- are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby

Tell your healthcare provider about all the medicines you take.



You should receive Evrysdi from the pharmacy as a liquid. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement



Avoid getting Evrysdi on your skin or in your eyes. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water

The most common side effects of Evrysdi include:



- For later-onset SMA:
 - fever
 - diarrhea
 - rash
- For infantile-onset SMA:
 - fever
 - diarrhea
 - rash
 - runny nose, sneezing and sore throat (upper respiratory infection)
 - lung infection (lower respiratory infection)
 - constipation
 - vomiting
 - cough

These are not all of the possible side effects of Evrysdi. For more information on the risk and benefits profile of Evrysdi, ask your healthcare provider or pharmacist.

You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at 1-888-835-2555.

Please see full Prescribing Information for additional Important Safety Information.